



Forward Looking Statement

This presentation and the oral commentary may contain forward-looking statements that involve future events. These forward-looking statements include terminology such as "believe," "will," "may," "estimate," "continue," "anticipate," "contemplate," "intend," "target," "project," "should," "plan," "expect," "predict," "could," "potentially" or the negative of these terms. All statements other than statements of historical facts contained in this presentation, including statements regarding our future financial condition, timing for and outcomes of clinical results, prospective products, preclinical and clinical pipelines, regulatory objectives, business strategy and plans and objectives for future operations, are forward looking statements Forward-looking statements are our current statements regarding our intentions, beliefs, projections, outlook, analyses or current expectations concerning, among other things, our anticipated significant milestones in 2021; the timing of our ongoing and planned clinical development; the sufficiency of our cash, cash equivalents and investments to fund our operations into the fourth quarter of 2023; our development programs for Zokinvy generally; and the potential approval of Zokinvy in jurisdictions outside of the U.S., including the EU in 2021; our progression and continued enrollment of our Phase 3 D-LIVR study in HDV; our ability to maintain supply of our commercial and clinical trial materials; our plans to advance Lambda in HDV in the U.S. and EU; our progression of Lambda for COVID-9 and Avexitide for PBH and HI; our ability to finance the continued advancement of our development pipeline products; and the potential for success of any of our product candidates. These statements concern product candidates that have not yet been approved for marketing by the U.S. Food and Drug Administration (FDA). No representation is made as to their safety or effectiveness for the purposes for which they are being investigated. Various important factors could cause actual results or events to differ materially from the forward-looking statements that Eiger makes, including additional applicable risks and uncertainties described in the "Risk Factors" sections in the Quarterly Report on Form 10-Q for the guarter ended March 31, 2021 and Eiger's subsequent filings with the SEC. The forward-looking statements contained in this press release are based on information currently available to Eiger and speak only as of the date on which they are made. Eiger does not undertake and specifically disclaims any obligation to update any forward-looking statements, whether as a result of any new information, future events, changed circumstances or otherwise. Additional information may be available in press releases or other public announcements and public filings made after the date of this presentation.

This presentation concerns certain products that have not yet been approved for marketing by the FDA. No representation is made as to their safety or effectiveness for the purposes for which they are being investigated.

© 2021 Eiger Biopharmaceuticals, Inc., all rights reserved. All trademarks belong to their respective owners.



Corporate Highlights

BUILDING A FULLY-INTEGRATED ORGANIZATION, SCALABLE FOR FUTURE GROWTH



- U.S. Commercial Launch: Progeria and processing-deficient progeroid laminopathies
- EMA approval expected in 2H 2021

HDV: >\$1B Opportunity

- LONAFARNIB: only oral therapy in Phase 3 D-LIVR study
- LAMBDA: a well-tolerated interferon entering Phase 3 LIMT-2 study

Lambda for COVID-19

- Positive Phase 2 results in newly diagnosed, non-hospitalized patients
- Phase 3 TOGETHER platform study enrolling in Brazil

Avexitide for Rare Metabolic Disorders

- PBH: Concurrence with FDA and EMA on single pivotal study
- HI: FDA Rare Pediatric Disease Designation; PRV eligible

Strong Cash Position

- Well-funded with ~\$160.5M cash as of 3/31/21
- Expected to fund planned operations into Q4 2023




Expected Key Milestones in 2021

CATALYSTS FOR VALUE CREATION













FDA Approved to Reduce the Risk of Mortality in capsules 50 mg/75 mg Hutchinson-Gilford Progeria Syndrome (Progeria)





*Used with permission



Progeria: Ultra-Rare, Fatal, Premature Aging Pediatric Disease



- Point mutation in the Lamin A gene
 - Results in a farnesylated aberrant protein, Progerin
 - Disruption of scaffold structure of the nuclear membrane
- Accelerated cardiovascular decline
- Average lifespan = 14.5 years without ZOKINVY treatment
- 90+ children and young adults treated with ZOKINVY
- ZOKINVY demonstrated survival benefit:
 - 60% reduction in risk of mortality
 - 2.5-year increase in mean survival
- Most common AEs are GI related





W/W Prevalence ~ 400 Children with Progeria



U.S. COMMERCIAL LAUNCH JANUARY 2021



Patients Identified in U.S.

- -FDA-approved
- -Commercial launch Jan 2021



Patients Identified in EU

-EMA approval expected 2H 2021







Eiger HDV Franchise



POTENTIAL FOR CONVENIENCE AND OPTIONALITY FOR HDV PATIENTS



- Potential HDV cure and maintenance therapies
- Foundational therapies for future combinations



HDV is Always a Co-infection with HBV

HDV REQUIRES HBsAg TO COMPLETE VIRUS ASSEMBLY


HDV HDV consists of HDV Large a single stranded, **HBsAg Acquired Through** genome delta antigen circular RNA virus, **PROTEIN PRENYLATION** with an envelope This is mechanism made up of HBsAg targeted by lonafarnib Small delta antigen **HBsAg HBV** surface antigen **HBV**



HDV: Most Severe Form of Viral Hepatitis



50% OF HDV-INFECTED PATIENTS ARE CIRRHOTIC AT DIAGNOSIS





12M HDV Patients Worldwide

~4-6% OF HBV-INFECTED POPULATION



Migration
Contributing to
Globalization
of Disease



\$1B+ HDV Market Opportunity



ONLY 3% MARKET PENETRATION REQUIRED











No FDA-Approved HDV Therapies: Treatments in Development

CHRONIC THERAPIES: CONVENIENCE MATTERS TO PATIENTS















Lonafarnib for HDV



FIRST AND ONLY ORAL AGENT IN DEVELOPMENT FOR HDV

- Well-characterized in patients
 - > 2,000 patients dosed in oncology program by Merck (Schering)
 - > 90 children dosed in Progeria program by Boston Children's Hospital
 - > 170 patients dosed in HDV program
 - Longest duration of dosing > 10 years
- Most common AEs are GI related
- Orphan Designation U.S. and EU
- FDA **Breakthrough Therapy** Designation
- EMA **PRIME** Designation
- Patent estate covers broad range of lonafarnib + ritonavir doses and durations











HDV-RNA REDUCTION IMPROVES CLINICAL OUTCOMES

Interferon- α for 48 Weeks with 15 year Follow Up

Change in HDV-RNA








Lonafarnib Phase 2 Data



TWO LONAFARNIB-BASED REGIMENS IDENTIFIED FOR REGISTRATION








D-LIVR Phase 3 Global Study



MULTIPLE PATHWAYS TO APPROVAL



Primary Endpoint at Week 48

≥ 2 log decline in HDV RNA

Normalization of ALT

Secondary Endpoint at Week 48

Histologic improvement Improvement of fibrosis

All patients will be maintained on background HBV nucleoside therapy. Superiority over PEG IFN-alfa-2a not required.



^{*} biopsy


Study Mirrors HDV Global Footprint



COMPLETION OF ENROLLMENT EXPECTED IN 2021

>75% Enrolled*
Full Enrollment in 2021

22 116
COUNTRIES SITES





Peginterferon Lambda (Lambda)


A WELL TOLERATED TYPE III INTERFERON




- Binds to a unique receptor vs type I IFN-a
 - Highly expressed on hepatocytes
 - Limited expression on hematopoietic and CNS cells
- Uses similar downstream signaling pathway to IFN-a
- 3,000+ patients in 19 clinical trials (HCV / HBV / HDV)
- Orphan Designation in U.S. and EU
- FDA Breakthrough Therapy Designation
- Composition of matter and method of use patents

Lambda Receptors Highly Expressed in the Liver

LAMBDA RECEPTORS NOT WIDELY DISTRIBUTED THROUGHOUT BODY



IFN- α RECEPTORS WIDELY DISTRIBUTED THROUGHOUT BODY








LVMT-1 Phase 2 Lambda Monotherapy Study Results

36% DURABLE VIROLOGIC RESPONSE (DVR) WITH LAMBDA






<u>L</u> <u>✓ MT – 2</u> Lambda Phase 3 Study



FDA & EMA CONCURRENCE ON SINGLE PIVOTAL STUDY






<u>L</u> ✓ MT – 2 Lambda Phase 3 Study



SINGLE PIVOTAL STUDY STARTING IN 2021



Primary Endpoint*


DVR at 24 Weeks Post-TRx (Arm 1) versus 12 Weeks Post-No TRx (Arm 2)



First-In-Class Treatments in Development for HDV



POTENTIAL FOR CONVENIENCE AND OPTIONALITY FOR HDV PATIENTS





Lambda for COVID-19



POTENTIAL AS A CONVENIENT, OUTPATIENT THERAPY FOR NEWLY DIAGNOSED PATIENTS

- Positive Phase 2 ILIAD results in newly diagnosed COVID-19 outpatients
- Potential to improve clinical outcomes and curb community spread
- Treatment well tolerated*
- Resistance due to variants of SARS-CoV-2 may not be an issue due to mechanism of action
- Provided in a prefilled syringe as a convenient, single dose, outpatient treatment
- Phase 3 TOGETHER investigator sponsored study enrolling in Brazil





Peginterferon lambda for the treatment of outpatients with COVID-19: a phase 2, placebo-controlled randomised trial



Jordan J Feld*, Christopher Kandel*, Mia J Biondi*, Robert A Kozak, Muhammad Atif Zahoor, Camille Lemieux, Sergio M Borgia, Andrea K Boggild, Jeff Powis, Janine McCready, Darrell H S Tan, Tiffany Chan, Bryan Coburn, Deepali Kumar, Atul Humar, Adrienne Chan, Braden O'Neil, Seham Noureldin, Joshua Booth, Rachel Hong, David Smookler, Wesam Aleyadeh, Anjali Patel, Bethany Barber, Julia Casey, Ryan Hiebert, Henna Mistry, Ingrid Choong, Colin Hislop, Deanna M Santer, D Lorne Tyrrell, Jeffrey S Glenn, Adam J Gehring, Harry L A Janssen, Bettina E Hansen

ILIAD Study Design

- Single dose Lambda vs placebo (1:1), N = 60
- Mild to moderate, non-hospitalized COVID-19 patients
- Primary endpoint: Proportion of patients with a negative SARS-CoV-2 nasopharyngeal swab at Day 7
- Follow-up through Day 14
- Mean baseline viral load = 6.71 log copies/mL





Peginterferon lambda for the treatment of outpatients with COVID-19: a phase 2, placebo-controlled randomised trial



Jordan J Feld*, Christopher Kandel*, Mia J Biondi*, Robert A Kozak, Muhammad Atif Zahoor, Camille Lemieux, Sergio M Borgia, Andrea K Boggild, Jeff Powis, Janine McCready, Darrell H S Tan, Tiffany Chan, Bryan Coburn, Deepali Kumar, Atul Humar, Adrienne Chan, Braden O'Neil, Seham Noureldin, Joshua Booth, Rachel Hong, David Smookler, Wesam Aleyadeh, Anjali Patel, Bethany Barber, Julia Casey, Ryan Hiebert, Henna Mistry, Ingrid Choong, Colin Hislop, Deanna M Santer, D Lorne Tyrrell, Jeffrey S Glenn, Adam J Gehring, Harry L A Janssen, Bettina E Hansen

ILIAD Study Results

- Lambda was over 4x more likely to clear infection within 7 days vs placebo
- For baseline viral load > 6 log copies/mL, 79% (Lambda) vs 38% (placebo) (p=0.012) clear by Day 7
- Fewer emergency room / hospital visits with Lambda (1) vs placebo (4)
- Lambda has potential to improve clinical outcomes and curb community spread
- Resistance due to variants of SARS-CoV-2 may not be an issue with Lambda due to mechanism of action
- Treatment well tolerated similar side effect profile to placebo









FIRST PATIENTS DOSED WITH LAMBDA IN JUNE 2021

- Ongoing, multi-center, investigator-sponsored, randomized, placebo-controlled Phase 3 study
- Evaluating multiple therapeutics in newly diagnosed, non-hospitalized patients with COVID-19
- TOGETHER now includes an investigational arm of Lambda as a single subcutaneous dose
- Lambda arm to enroll up to 800 high-risk, non-hospitalized patients
- Primary endpoint is reduction of emergency room visits and hospitalizations
- Protocol includes interim analysis for futility
- Recruiting at eleven sites in Brazil and expanding to include a site in Toronto, Canada



together • COVID-19 Phase 3 Platform Study

PRINCIPAL INVESTIGATORS



EDWARD MILLS, PhD



- Principal investigator, TOGETHER COVID-19 trial
- Professor of Health Research
 Methods, Evidence, and
 Impact at McMaster University,
 Vancouver, Canada



GILMAR REIS, MD, PhD




- Co-investigator, TOGETHER COVID-19 trial
- Director of the Outpatient
 Research Clinic at Cardresearch
- Associate Professor of Medicine at Pontifical Catholic University of Minas Gerais, Brazil



Avexitide Is a First-in-Class GLP-1 Antagonist

31 AMINO ACID FRAGMENT OF BYETTA (EXENATIDE), A GLP-1 AGONIST

- Novel Liquid Formulation Developed
- Sub-cutaneous delivery
- Targeted MOA for PBH and CHI
- Differential dose/device strategies for PBH & HI
- Patent protection will provide market exclusivity through at least 2039





Avexitide for PBH

PHASE 3 READY IN 2022

POST-BARIATRIC HYPOGLYCEMIA (PBH)





- Complication of bariatric surgery
- Dangerously low blood sugar after meals
- ~5-10% of Roux-en-Y Gastric Bypass
- ~2.5% of Vertical Sleeve Gastrectomy
- ~150K PBH patients in U.S. & EU5
- Other procedures: esophagectomy, gastrectomy, Nissen fundoplication

PBH results in **SEVERE HYPOGLYCEMIA**: altered mental status, loss of consciousness, seizures, coma

CLINICAL & REGULATORY STATUS

- Proof of concept demonstrated in 54 patients dosed across
 4 completed Phase 2 studies
- Most recent Phase 2 data published in JCEM 2021
- FDA Breakthrough Therapy Designation for Hypoglycemic Hypoglycemia
- Concurrence with FDA and EMA on single pivotal study
 (N=90) with end points previously demonstrated in Phase 2

Phase 3 Ready in 2022:
Manufacturing and device development ongoing



Avexitide for HI



FDA DISCUSSIONS ONGOING FOR PHASE 3 STUDY DESIGN

CONGENITAL HYPERINSULINISM (HI)



- Ultra-rare pediatric metabolic disorder
- Most frequent cause of persistent hypoglycemia in neonates and children
- >3,300 patients in U.S. & EU5
- Well-defined market; multiple therapies in development
- Near-total pancreatectomy is indicated

CLINICAL & REGULATORY STATUS

- Proof of concept demonstrated in 39 patients across 3
 Phase 2 studies (neonates, children, adolescents/adults)
- Clinically significant endpoint (glucose infusion rate) demonstrated
- FDA Rare Pediatric Disease Designation; PRV eligible
- Differentiated from PBH based on dose and device

HI results in PERMANENT BRAIN DAMAGE with neurodevelopmental deficits in up to 50% of patients

FDA discussions ongoing for Phase 3 neonate/child studies



Diverse, Late-Stage Pipeline

TARGETED INDICATION		DRUG	ORPHAN US / EU	BREAKTHROUGH THERAPY	RARE PEDIATRIC DISEASE	STATUS & UPCOMING MILESTONES
¥	Progeria and Progeroid Laminopathies	Zokínvy [®] (lonafarnib) capsules 50 mg/75 mg			PRV Sold	FDA APPROVED; EMA Approval 2H21
	Hepatitis Delta Virus	Lonafarnib			N/A	Phase 3 Enrollment Completion in 2021
		Peginterferon Lambda			N/A	Phase 3 Start in 2021
	COVID-19		N/A	N/A	N/A	Phase 3 Enrolling
7	Post-Bariatric Hypoglycemia	Avexitide	✓	~	N/A	Phase 3 Ready
श्रीह	Congenital Hyperinsulinism					Phase 2





- Phase 3 *D-LIVR* full enrollment by end of 2021
- Phase 3 *LIMT-2* Lambda study initiation in 2H 2021
- Zokinvy® U.S. launch and EMA approval in 2021
- Avexitide manufacturing and device development in 2021
- Phase 3 TOGETHER study of Lambda for COVID-19 enrolling
- Strong cash balance: planned operations funded into 4Q 2023

